



The Disorder Named AD/HD - CHADD Fact Sheet #1

Occasionally, we may all have difficulty sitting still, paying attention or controlling impulsive behavior. For some people, the problems are so pervasive and persistent that they interfere with their lives, including home, academic, social and work settings.

Attention-deficit/hyperactivity disorder (AD/HD) is characterized by developmentally inappropriate levels of inattention, impulsivity, and hyperactivity. AD/HD is a neurobiological disorder that affects 3 to 7 percent of school-age children.^{1,2,3,4} Until relatively recently, it was believed that children outgrew AD/HD in adolescence as hyperactivity often diminishes during the teen years. However, it is now known that AD/HD nearly always persists from childhood through adolescence and that many symptoms continue into adulthood. In fact, current research reflects rates of roughly 2 to 4 percent among adults.⁵

Although individuals with this disorder can be very successful in life, without identification and proper treatment, AD/HD may have serious consequences, including school failure, family stress and disruption, depression, problems with relationships, substance abuse, delinquency, risk for accidental injuries and job failure. Early identification and treatment are extremely important.

Medical science first documented children exhibiting inattentiveness, impulsivity and hyperactivity in 1902. Since that time, the disorder has been given numerous names, including minimal brain dysfunction, hyperkinetic reaction of childhood and attention-deficit disorder with or without hyperactivity. With the *Diagnostic and Statistical Manual, fourth edition (DSM-IV)* classification system, the disorder has been renamed attention-deficit/hyperactivity disorder, or AD/HD. The current name reflects the importance of the inattention characteristics of the disorder as well as the other characteristics of the disorder, such as hyperactivity and impulsivity.

The Symptoms

Typically, AD/HD symptoms arise in early childhood, unless associated with some type of brain injury later in life. Some symptoms persist into adulthood and may pose life-long challenges. Although the official diagnostic criteria state that the onset of symptoms must occur before age seven, leading researchers in the field of AD/HD argue that criterion should be broadened to include onset anytime during childhood⁶. The symptom-related criteria for the three primary subtypes are adapted from *DSM-IV* and summarized as follows:

AD/HD predominantly inattentive type: (AD/HD-I)

- Fails to give close attention to details or makes careless mistakes.
- Has difficulty sustaining attention.
- Does not appear to listen.

- Struggles to follow through on instructions.
- Has difficulty with organization.
- Avoids or dislikes tasks requiring sustained mental effort.
- Loses things.
- Is easily distracted.
- Is forgetful in daily activities.

AD/HD predominantly hyperactive-impulsive type: (AD/HD-HI)

- Fidgets with hands or feet or squirms in chair.
- Has difficulty remaining seated.
- Runs about or climbs excessively.
- Difficulty engaging in activities quietly.
- Acts as if driven by a motor.
- Talks excessively.
- Blurts out answers before questions have been completed.
- Difficulty waiting or taking turns.
- Interrupts or intrudes upon others.

AD/HD combined type: (AD/HD-C)

- Individual meets both sets of inattention and hyperactive/impulsive criteria.

Youngsters with AD/HD often experience delays in independent functioning and may therefore behave in ways more like younger children.⁶ In addition, AD/HD frequently co-occurs with other conditions, such as depression, anxiety or learning disabilities. For example, in 1999, NIMH research indicated that two-thirds of children with AD/HD have a least one other co-existing condition.⁷ When co-existing conditions are present, academic and behavioral problems, as well as emotional issues, may be more complex.

Teens with AD/HD present a special challenge. During these years, academic and organizational demands increase. In addition, these impulsive youngsters are facing typical adolescent issues: discovering their identity, establishing independence, dealing with peer pressure, exposure to illegal drugs, emerging sexuality, and the challenges of teen driving.

Recently, deficits in executive function have emerged as key factors impacting academic and career success⁸. Simply stated, executive function refers to the “variety of functions within the brain that activate, organize, integrate and manage other functions.”⁹ This permits individuals to appreciate the longer-term consequences of their actions and guide their behavior across time more effectively.¹⁰ Critical concerns include deficits in working memory and the ability to plan for the future, as well as maintaining and shifting strategies in the service of long-term goals.

The Diagnosis

Determining if a child has AD/HD is a multifaceted process. Many biological and psychological problems can contribute to symptoms similar to those exhibited by children with AD/HD. For example, anxiety, depression and certain types of learning disabilities may cause similar symptoms. In some cases, these other conditions may actually be the primary diagnosis; in others, these conditions may co-exist with AD/HD.

There is no single test to diagnose AD/HD. Therefore, a comprehensive evaluation is necessary to establish a diagnosis, rule out other causes and determine the presence or absence of co-existing conditions. Such an evaluation requires time and effort and should include a careful history and a clinical assessment of the individual's academic, social, and emotional functioning and developmental level. A careful history should be taken from the parents and teachers, as well as the child, when appropriate. Checklists for rating AD/HD symptoms and ruling out other disabilities are often used by clinicians; these age-normed instruments help to ensure that the symptoms are extreme for the child's developmental level.

There are several types of professionals who can diagnose AD/HD, including school psychologists, clinical psychologists, clinical social workers, nurse practitioners, neurologists, psychiatrists and pediatricians. Regardless of who does the evaluation, the use of the *Diagnostic and Statistical Manual IV* diagnostic criteria for AD/HD is necessary. A medical exam by a physician is important and should include a thorough physical examination, including assessment of hearing and vision, to rule out other medical problems that may be causing symptoms similar to AD/HD. In rare cases, persons with AD/HD also may have a thyroid dysfunction. Only medical doctors can prescribe medication if it is needed. Diagnosing AD/HD in an adult requires an evaluation of the history of childhood problems in behavior and academic domains, as well as examination of current symptoms and copying strategies. For more information, read the information and resource sheet on "The Diagnosis of AD/HD in Adults."

The Causes

Multiple studies have been conducted to discover the cause of the disorder. Research clearly indicates that AD/HD tends to run in families and that the patterns of transmission are to a large extent genetic.^{11,12} More than 20 genetic studies, in fact, have shown evidence that AD/HD is strongly inherited. Yet AD/HD is a complex disorder, which is undoubtedly the result of multiple interacting genes. Other causal factors (such as low birthweight, prenatal maternal smoking, and additional prenatal problems) may contribute to other cases of AD/HD.¹³⁻¹⁶ Problems in parenting or parenting styles may make AD/HD better or worse, but these do not cause the disorder. AD/HD is clearly a brain-based disorder. Currently research is underway to better define the areas and pathways that are involved.

Prognosis and Long-term Outcomes

Children with AD/HD are at risk for potentially serious problems in adolescence: academic underachievement and school failure, problems in social relations, risk for antisocial behavior patterns, teen pregnancy, and adverse driving consequences.¹⁷ As noted above, AD/HD persists from childhood to adolescence in the vast majority of cases, although the symptom area of motor activity tends to diminish with time. Furthermore, up to two-thirds of children with AD/HD continue to experience significant symptoms in adulthood. Yet many adults with AD/HD learn coping strategies and compensate quite well.^{18,19} A key to good outcome is early identification and treatment.

Multimodal Treatment

AD/HD in children often requires a comprehensive approach to treatment called "*multimodal*" and includes:

- Parent and child education about diagnosis and treatment
- Specific behavior management techniques
- Medication
- Appropriate school programming and supports

Treatment should be tailored to the unique needs of each child and family. Research from the landmark NIMH Multimodal Treatment Study of AD/HD is very encouraging.²⁰ Children who received carefully monitored medication, alone or in combination with behavioral treatment, showed significant improvement in their behavior at home and school plus better relationships with their classmates and family than did children receiving lower quality care.

Psychostimulants are the most widely used class of medication for the management of AD/HD related symptoms. Approximately 70 to 80 percent of children with AD/HD respond positively to psychostimulant medications.²¹ Significant academic improvement is shown by students who take these medications: *increases in* attention and concentration, compliance and effort on tasks, as well as amount and accuracy of schoolwork, plus *decreased* activity levels, impulsivity, negative behaviors in social interactions and physical and verbal hostility.²² A new, nonstimulant medication—atomoxetine--appears to have similar effects as the stimulants.

Other medications that may decrease impulsivity, hyperactivity and aggression include some antidepressants and antihypertensives. However, each family must weigh the pros and cons of taking medication (see the CHADD fact sheet, “Evidence-based Medication Management for Children and Adolescents with AD/HD”).

Behavioral interventions are also a major component of treatment for children who have AD/HD. Important strategies include being consistent and using positive reinforcement, and teaching problem-solving, communication, and self-advocacy skills. Children, especially teenagers, should be actively involved as respected members of the school planning and treatment teams (see the CHADD fact sheet, “Evidence-based Psychosocial Treatment for Children and Adolescents with AD/HD”).

School success may require a variety of classroom accommodations and behavioral interventions. Most children with AD/HD can be taught in the regular classroom with minor adjustments to the environment. Some children may require special education services if an educational need is indicated. These services may be provided within the regular education classroom or may require a special placement outside of the regular classroom that meets the child’s unique learning needs (see the CHADD fact sheet on educational rights).

Adults with AD/HD may benefit from learning to structure their environment. In addition, medications effective for childhood AD/HD are also helpful for adults who have AD/HD. While little research has been done on interventions for adults, diagnosis and treatment are still important.

Summary

Although the symptoms of AD/HD—inattention, impulsivity and hyperactivity—are present to some extent in most children, when these symptoms are developmentally extreme, pervasive and persistent a diagnosis of AD/HD is warranted. This diagnostic category is associated with significant impairment in family relations, peer interactions, school achievement, and risk for accidental injury, which are domains of crucial importance for healthy and successful development. Because AD/HD can become a lifelong disorder, careful diagnosis and treatment are essential. CHADD is seeking out solutions that will lead to improved quality of life for children, adolescents and adults.

Suggested reading

Barkley, R. (1998). *Attention Deficit Hyperactivity Disorders: A Handbook for Diagnosis and Treatment*. New York: Guilford Press.

Brown, T.E. (2000). *Attention-deficit Disorders and Comorbidities in Children, Adolescents, and Adults*. Washington, D.C.: American Psychiatric Press, Inc.

Dendy, C.A.Z and Ziegler, Alex. (2003). *A Bird's-Eye View of Life with ADD and ADHD: Advice from Young Survivors*. Cedar Bluff, AL: Cherish the Children.

DuPaul, G.J. & Stoner, G. (2003). *ADHD in the Schools: Assessment and Intervention Strategies*. New York, NY: The Guilford Press.

Goldstein, S. (1998). *Managing Attention Deficit Hyperactivity Disorder in Children: A Guide for Practitioners*. New York, NY: John Wiley & Sons.

Hallowell, E.M. and Ratey, J.J. (1995). *Driven to Distraction: Recognizing and Coping with Attention Deficit Disorder from Childhood through Adulthood*. New York: Simon & Schuster.

Ingersoll, Barbara D. (1995). *Distant Drums, Different Drummers: A Guide for Young People with ADHD*. Germantown, MD: Cape Publications.

Jensen, P.S. and Cooper, J.R., editors. (2002) *Attention Deficit Hyperactivity Disorder: State of Science--Best Practices*. Kingston, NJ: Civic Research Institute.

Jensen, P. (2004) *Making the System Work for Your Child with ADHD: An Expert Parent's Guide to Getting the Best Care*. New York, NY: Guilford Press.

Jones, Clare. (2003) *Practical Suggestions for ADHD*. East Moline, IL: LinguiSystems Publications.

Nadeau, Kathleen G. and Quinn, Patricia O., editors. (2002) *Understanding Women with AD/HD*. Silver Spring, MD: Advantage Books.

Nadeau, Kathleen G.; Littman, Ellen B.; and Quinn, Patricia O. (1999) *Understanding Girls With AD/HD*. Silver Spring, MD: Advantage Books.

Parker, H.C. (2002). *Problem Solver Guide for Students with ADHD: Ready-to-Use Interventions for Elementary and Secondary Students with Attention Deficit Hyperactivity Disorder*. Plantation, FL: Impact Publications.

Rief, S. (2003). *The AD/HD Book of Lists*. San Francisco, CA: Jossey-Bass.

Robin, A.L. (1998). *ADHD in Adolescents: Diagnosis and Treatment*. New York, NY: The Guilford Press.

Solden, Sari. (1995). *Women with Attention Deficit Disorder: Embracing disorganization at Home and in the Workplace*. Grass Valley, CA: Underwood Books.

Weiss, Lynn. (1997). *Attention Deficit Disorder in Adults: Practical Help and Understanding*. Lanham, MD: Taylor Trade Publishing.

Wilens, Timothy (1999). *Straight Talk about Psychiatric Medications for Kids*. New York, NY: Guilford Press.

References

1. American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders: DSM-IV* (4th ed., text revision), Washington, D.C.: American Psychiatric Association.
2. Mayo Clinic. (2002). How Common is Attention-Deficit/Hyperactivity Disorder? *Archives of Pediatrics and Adolescent Medicine* 156(3): 209-210.
3. Mayo Clinic. (2001). Utilization and Costs of Medical Care for Children and Adolescents with and without Attention-Deficit/Hyperactivity Disorder, *Journal of the American Medical Association* 285(1): 60-66.
4. Surgeon General of the United States. (1999) *Mental Health: A Report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services.
5. Murphy, K. R., & Barkley, R.A. (1996) The prevalence of DSM-IV symptoms of AD/HD in adult licensed drivers: Implications for clinical diagnosis. *Comprehensive Psychiatry*, 37, 393-401.
6. Barkley, RA. (1998). *Attention deficit hyperactivity disorders: A handbook for diagnosis and treatment*. New York: Guilford Press.
7. MTA Cooperative Group. (1999) A 14-month randomized clinical trial of treatment strategies for attention deficit hyperactivity disorder. *Archives of General Psychiatry*, 56, 12.
8. Barkley, RA. (1998). *Attention deficit hyperactivity disorders: A handbook for diagnosis and treatment*. New York: Guilford Press.
9. Brown, T.E. (2000). *Attention-deficit Disorders and Comorbidities in Children, Adolescents, and Adults*. Washington, D.C.: American Psychiatric Press, Inc.
10. Fuster, J.M. (1997). *The prefrontal cortex: anatomy, physiology, and neuropsychology of the frontal lobe*. Philadelphia: Lippincott-Raven.
11. Tannock, R (1998). Attention deficit hyperactivity disorder: Advances in cognitive, neurobiological, and genetic research. *Journal of Child Psychology and Psychiatry*, 39, 65-99.
12. Swanson, JM, and Castellanos, FX (2002). Biological Bases of ADHD—Neuroanatomy, Genetics, and Pathophysiology. In P.S. Jensen and J.R. Cooper (eds). *Attention deficit hyperactivity disorder: State of the science, best practices*, pp. 7-1—7-20. Kingston, New Jersey.
13. Connor, D.R. (2002). Preschool Attention deficit hyperactivity disorder: A review of prevalence, diagnosis, neurobiology, and stimulant treatment. *Journal of Developmental Behavior Pediatrics* 23 (1Suppl): S1-S9.
14. Wilens, T.E., Biederman, J.; Brown, S.; Tanguay, S.; Monteaux, M.C.; Blake, C.; Spencer, J.J. (2002). Psychiatric co-morbidity and functioning in clinically referred preschool children and school age youths with AD/HD. *Journal of the American Academy of Child and Adolescent Psychiatry* 4(3): 26-28.
15. Teeter, P. (1998). *Interventions for AD/HD*. New York: Guilford Press.
16. Jones, C. (2003). *Practical Suggestions for AD/HD*. East Moline, IL: Linguist-Systems.
17. Barkley, RA. (1998). *Attention deficit hyperactivity disorders: A handbook for diagnosis and treatment*. New York: Guilford Press.
18. Barkley, RA, Fischer, M., Fletcher, K., & Smallish, L. (2001) *Young adult outcome of hyperactive children as a function of severity of childhood conduct problems, I: Psychiatric status and mental health treatment*. Submitted for publication.

-
19. Weiss G, Hechtman, L, Milroy T et al. (1985). Psychiatric studies of hyperactives as adults: a controlled prospective 15-year follow-up of 63 hyperactive children. *Journal of the American Academy of Child Psychiatry*, 23, 211-220.
 20. MTA Cooperative Group. (1999) A 14-month randomized clinical trial of treatment strategies for attention deficit hyperactivity disorder. *Archives of General Psychiatry*, 56, 12.
 21. Spencer, T., Wilens, T., Biederman, J., Faraone, S. V., Ablon, J. S., & Lapey, K. (1995). A double-blind, crossover comparison of methylphenidate and placebo in adults with childhood-onset attention-deficit hyperactivity disorder. *Archives of General Psychiatry*, 52, 434-443.
 22. Swanson, JM, McBurnett K, et al (1993) Effect of stimulant medication on children with attention deficit disorder: a “review of reviews.” *Exceptional Children*, 60, 154-162.

© 2005 Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD). All rights reserved. Reproduction in whole or in part without written permission from CHADD is prohibited.

For further information about AD/HD or CHADD, please contact:

**CHADD National Resource Center on AD/HD
8181 Professional Place, Suite 150, Landover, MD 20785
1-800-233-4050
www.help4adhd.org**

Please also visit the CHADD Web site at www.chadd.org